arXiv:1212.5302v1 [math.RT] 20 Dec 2012

IRREDUCIBILITY CRITERION FOR REPRESENTATIONS INDUCED BY ESSENTIALLY UNITARY ONES (CASE OF NON-ARCHIMEDEAN $GL(n,\mathcal{A})$)

MARKO TADIĆ

Abstract. The fundamental representations for the description of the unitary dual of $GL(n)$ over a local field F are the Speh representations, which are denoted by $u(\delta,m)$. Let $\mathcal{A}$ be a finite dimensional central division algebra over F. One can define in a similar way representations $u(\delta,m)$ for $GL(n,\mathcal{A})$, and they play a similar fundamental role for the unitary duals of these groups. In the non-archimedean case, for two such representations $u(\delta_1,m_1)$ and $u(\delta_2,m_2)$ of $GL(n_1,\mathcal{A})$ and $GL(n_2,\mathcal{A})$ respectively, and for $\alpha_1,\alpha_2\in\mathbb{R}$, we obtain a necessary and sufficient condition that the parabolically induced representation

$$\mathrm{Ind}^{GL(n_1+n_2,\mathcal{A})}(|\ |_F^{\alpha_1}u(\delta_1,m_1)\otimes|\ |_F^{\alpha_2}u(\delta_2,m_2))$$

is irreducible. Our motivation for working on this problem comes from our work on unitary duals of classical (symplectic, orthogonal and unitary) groups, related to constructions of complementary series for these groups.

Contents

1. Introduction 2
2. Notation 7
3. Some criteria for reducibility and irreducibility 15
4. Contacts of non-induced essentially unitarizable representations 17
5. Linked case 20
6. Non-linked case 23
7. Reducibility criterion 33

Date: December 24, 2012.

2000 *Mathematics Subject Classification.* Primary: 22E50.

Key words and phrases. non-archimedean local fields, division algebras, general linear groups, Speh representations, parabolically induced representations, reducibility, unitarizability.

The author was partly supported by Croatian Ministry of Science, Education and Sports grant #037-0372794-2804.

8. Relation with a result of C. Mœglin and J.-L. Waldspurger 35
References 36

1. Introduction

The investigations of this paper are entirely for non-archimedean general linear groups, but the main motivation for them comes from the classical groups (symplectic, orthogonal and unitary ones). We explain the motivation, considering the case of symplectic groups.

Let F be a local non-archimedean field and $Sp(2n, F)$ the symplectic group of $2n \times 2n$ matrices. We consider the problem of unitarizability for these groups. In [28] we have reduced the problem of unitarizability to the case of so called weakly real representations. Roughly, for an irreducible representation π, this means that the unitary parts of irreducible cuspidal representations of general linear groups that occur in description of π by parabolic induction are selfdual, i.e., isomorphic to their contragredients.

A significant part of irreducible unitary representations of classical groups consists of complementary series (see [13] or [16], for example). Thanks to [28], it is enough to take care of only weakly real complementary series. We try to give idea what (at least some of) these representations could look like. Let σ be an irreducible unitarizable representation of $Sp(2q, F)$ and $\pi_1, \ldots, \pi_l$ be irreducible selfdual representations of $GL(p_1, F), \ldots, GL(p_l, F)$. Let $n = p_1 + \cdots + p_l + q$. Suppose that we have a connected subset $X \subseteq \mathbb{R}^l$ such that parabolically the induced representations

$$\mathrm{Ind}_P^{Sp(2n,F)}(|\ |_F^{\alpha_1}\pi_1 \otimes \ldots \otimes |\ |_F^{\alpha_l}\pi_l \otimes \sigma), \quad (\alpha_1, \ldots, a_l) \in X, \tag{1.1}$$

are irreducible, where P is the appropriate parabolic subgroup (whose Levi factor is isomorphic to the direct product $GL(p_1, F) \times \ldots \times GL(p_l, F) \times Sp(2q, F)$). Suppose that for some particular $(\alpha_1, \ldots, \alpha_l) \in X$, the corresponding induced representation in (1.1) is unitarizable (in general, one gets such unitarizability using unitary parabolic induction from a proper parabolic subgroup, which may not always be associate of P). Then all the representations in (1.1) are unitarizable. Such sets of representations are called complementary series. So, the principal problem is to get a family of induced representations as above, containing a unitarizable one. This can be split into several problems. One of them we discuss below.

For a fixed representation from (1.1), all the semisimplifications of representations

$$\mathrm{Ind}_P^{Sp(2n,F)}(|\ |_F^{\pm\alpha_{\tau(1)}}\pi_{\tau(1)} \otimes \ldots \otimes |\ |_F^{\pm\alpha_{\tau(l)}}\pi_{\tau(l)} \otimes \sigma) \tag{1.2}$$

are the same, for all the permutations τ of $\{1, \ldots, l\}$ (see [24]). Therefore

$$\mathrm{Ind}_P^{Sp(2n,F)}(|\ |_F^{\alpha_1}\pi_1 \otimes \ldots \otimes |\ |_F^{\alpha_l}\pi_l \otimes \sigma)$$

is irreducible if and only if all the representations in (1.2) are irreducible. Now induction in stages and the requirement that representations in (1.1) are irreducible imply that all the representations

$$\mathrm{Ind}_{P'}^{GL(p_i+p_j,F)}(|\ |_F^{\alpha_i}\pi_i \otimes |\ |_F^{\pm\alpha_j}\pi_j), \quad 1 \leq i < j \leq l, \tag{1.3}$$

and

$$\mathrm{Ind}_{P''}^{Sp(2(p_i+q),F)}(|\ |_F^{\alpha_i}\pi_i \otimes \sigma), \quad 1 \leq i \leq l, \tag{1.4}$$

are irreducible when $(\alpha_1, \ldots, a_l) \in X$.

Therefore, for construction of general complementary series (1.1) of classical groups, we can start from an easier problem. We can first examine the irreducibility of representations (1.3) and (1.4) (for both of them, irreducibility must hold), and then try to see if this implies the irreducibility of representations from (1.1) (this is, for example, the case if all π_i and σ are tempered representations). For studying these two irreducibility questions (which are very hard for general irreducible representations), it would be very important to know for which representations π_i and σ we actually need to consider these irreducibility questions (see [13] and [16] for the case of generic and unramified duals).

Although the classifications in [21], [13] and [16] give some idea of the shape of the classification of unitary duals of classical groups that we can expect in general, serious work is still needed to make the expectation more explicit. Regarding complementary series, we do not know precisely which representations σ's we need to consider in (1.1). This is one of the most important problems (for the generic unitary dual these were generic irreducible tempered representations, while for the unramified unitary dual, these were irreducible unramified negative representations). Regarding representations π_i in (1.1), [21], [13] and [16] suggest that for these representations, natural candidates are the Speh representations

$$u(\delta, n),$$

which will be defined below[1].

In this paper we study the reducibility of representations (1.3) in the case of Speh representations, and more generally for general linear groups over central finite dimensional division algebras over F, and for the corresponding representations there. We get a necessary and sufficient criterion for the irreducibility of these representations.

Let us note that this problem was studied in the field case by C. Mœglin and J.-L. Waldspurger. They have obtained a (sufficient) irreducibility criterion for representations (1.3) in the case of Speh representations ([15], Lemma I.6.3). Since for the unitarizability, it is crucial to know if the sets in (1.1) are maximal (with given properties), it is important to us to know if their criterion is also necessary. Our paper shows that this is the case (for split general linear groups, which they consider). Actually, this is relatively easy to show (see the fifth section of this paper). Moreover, in the case of reducibility, we can determine two non-equivalent irreducible subquotients[2] (but we do not go into explicit formulas for them in this paper).

---

[1] Since we are considering the weakly real case, δ needs to be selfdual.

The proof of irreducibility in [15] is based on analytic properties of standard integral intertwining operators, normalized by L-functions and ϵ-factors. For a long time, we were thinking of proving the irreducibility result of [15] by different methods, basically in the scope of the Bernstein-Zelevinsky theory (since such irreducibility is a natural problem from that theory), and in this way get a different understanding of the result of C. Mœglin and J.-L. Waldspurger. This was another motivation for writing this paper.

The methods that we use in this paper are rather elementary. The principal tool in our paper is the combinatorial algorithm of C. Mœglin and J.-L. Waldspurger from [14] for the Zelevinsky involution. The starting point of our paper was the paper [5] of I. Badulescu, E. Lapid and A. Míngues. Our paper uses the general criterion for irreducibility obtained there. This criterion is not sufficient for the irreducibility in general, not even for the case that we consider. In the case where the criterion of [5] does not work, we apply a simple criterion of I. Badulescu from [3], based on the combinatorial algorithm of C. Mœglin and J.-L. Waldspurger (which is in general not necessary for the irreducibility, again even for the case that we consider).

Before we describe the reducibility criterion that we obtain in this paper, let us note that in the case of general complementary series of classical groups, we can have a much different situation than we have in [13] and [16], where the exponents in all complementary series were at most 1. The reason for the difference is that in the non-generic case, we can have cuspidal reducibilities as big as we want (see [29]).

In our forthcoming work, we plan to address some of the further questions related to the general complementary series of classical groups.

Let F be a local non-archimedean field and let $\mathcal{A}$ be a finite dimensional central division algebra of rank $d_{\mathcal{A}}$ over F. Put

$$G_p = GL(p, \mathcal{A}).$$

For an irreducible essentially square integrable representation δ of G_p, denote by s_δ the smallest positive real number such that

$$\mathrm{Ind}^{G_{2p}}(\delta \otimes |\ |_F^{s(\delta)}\delta)$$

reduces. Then $s_\delta \in \mathbb{Z}$ and $s_\delta | d_{\mathcal{A}}$. Let

$$\nu_\delta := |\ |_F^{s(\delta)}.$$

---

[2] By the same method we determined two irreducible subquotients at the end of complementary series in [22]. Clearly, if we have two non-equivalent irreducible subquotients, this implies the reducibility. This is the way we prove reducibility in this paper, using the Zelevinsky involution and the combinatorial algorithm of C. Mœglin and J.-L. Waldspurger from [14] for this involution.

Let ρ be an irreducible cuspidal representation of G_p and $n \in \mathbb{Z}_{\geq 0}$. Then we denote

$$[\rho, \nu_\delta^n \rho] := \{\rho, \nu_\delta \rho, \dots, \nu_\delta^n \rho\}.$$

The set $\Delta = [\rho, \nu_\delta^n \rho]$ is called a segment in cuspidal representations. For $a, b \in \mathbb{Z}$, the set $\{x \in \mathbb{Z}; a \leq x \leq b\}$ is called a $\mathbb{Z}$-segment. It is denoted by

$$[a, b]_{\mathbb{Z}}.$$

Let $\Delta = [\rho, \nu_\delta^n \rho]$. Consider the representation

$$\mathrm{Ind}^{G_{(n+1)p}}(\nu_\delta^n \rho \otimes \nu_\delta^{n-1} \rho \otimes \ldots \otimes \rho),$$

parabolically induced from the appropriate standard parabolic subgroup (see the second section). Then the above representation has a unique irreducible subrepresentation, denoted by

$$\delta(\Delta).$$

Further, $\delta(\Delta)$ is essentially square integrable, and we get all the essentially square integrable representations in this way.

Let δ be an irreducible essentially square integrable representation of G_p. Then the representation

$$\mathrm{Ind}^{G_{np}}(\nu_\delta^{(n-1)/2} \delta \otimes \nu_\delta^{(n-1)/2-1} \delta \otimes \ldots \otimes \nu_\delta^{-(n-1)/2} \delta),$$

parabolically induced from the appropriate standard parabolic subgroup, has a unique irreducible quotient, denoted by

$$u_{ess}(\delta, n).$$

It is essentially unitarizable[3] (i.e., becomes unitarizable after twist by a character; see [6]).

A different parameterization of representations $u(\delta, n)$ will be convenient for us. Let

$$\delta = \delta([\nu_\delta^b \rho, \nu_\delta^e \rho]).$$

Let

$$A = b - (n-1)/2, \qquad B = e - (n-1)/2,$$
$$C = b + (n-1)/2, \qquad D = e + (n-1)/2.$$

Then we denote

$$u_{ess}\begin{pmatrix} A & & B & \\ & C & & D \end{pmatrix}^{(\rho)} := u_{ess}(\delta([\nu_\delta^b \rho, \nu_\delta^e \rho]), n).$$

---

[3] It is not clear to us if it is appropriate to call these essentially unitarizable representations Speh representation (as they were called in [3] and [5], for example). Namely, by [2] and [8], these representations are usually not automorphic, and then do not have global counterparts, so lack a principal property of the Speh representation from the field case. The local representations, denoted by $\bar{u}(\delta, m)$ in [6] (with global counterparts), might be more appropriate to call Speh representations in this setting.

This notation is very natural from the point of Langlands classification (see the seventh section[4]).

We write

$$\begin{pmatrix} A & & B & \\ & C & & D \end{pmatrix} <_{strong} \begin{pmatrix} A' & & B' & \\ & C' & & D' \end{pmatrix} \iff A < A', B < B', C < C', D < D'.$$

Theorem 1.1. *Let* δ_1 *and* δ_2 *be irreducible essentially square integrable representations of* G_{p_1} *and* G_{p_2} *respectively, and* n_1, n_2 *positive integers. If the representation*

$$Ind^{G_{n_1p_1+n_2p_2}}(u_{ess}(\delta_1, n_1) \otimes u_{ess}(\delta_2, n_2)) \tag{1.5}$$

is reducible, then we can find an irreducible cuspidal representation ρ *and* $A_i, B_i, C_i, D_i \in \mathbb{Z}$, *such that*

$$u_{ess}(\delta_i, n_i) = u_{ess}\begin{pmatrix} A_i & & B_i & \\ & C_i & & D_i \end{pmatrix}^{(\rho)}$$

for $i = 1, 2$. *Now* (1.5) *reduces if and only if*

$$[A_1, D_1]_{\mathbb{Z}} \cup [A_2, D_2]_{\mathbb{Z}}$$

is a $\mathbb{Z}$*-segment, and the following holds:*

$$\begin{pmatrix} A_1 & & B_1 & \\ & C_1 & & D_1 \end{pmatrix} <_{strong} \begin{pmatrix} A_2 & & B_2 & \\ & C_2 & & D_2 \end{pmatrix} \quad or \quad \begin{pmatrix} A_2 & & B_2 & \\ & C_2 & & D_2 \end{pmatrix} <_{strong} \begin{pmatrix} A_1 & & B_1 & \\ & C_1 & & D_1 \end{pmatrix}.$$

A different formulation of the above theorem, in terms of segments defining the representations, can be found in the seventh section. Let us recall that [15] contains a stronger irreducibility result then the above theorem (for the field case). There the irreducibility of representations parabolically induced by tensor products of several essentially Speh representations which satisfy the assumptions for irreducibility of the above theorem is proved (see Proposition I.9 of [15]).

A puzzling question which arises is the following: does the criterion that we have proved in this paper generalize in a natural way to the ladder representation of [12]? The formulation of the criterion in Theorem 7.1 might be a better starting point for this question, than the above theorem.

Discussions with I. Badulescu, E. Lapid and C. Moeglin were helpful in the course of preparation of this paper. C. Jantzen's numerous corrections helped us a lot to improve the style of the paper. We are thankful to all them.

---

[4] The representation $u_{ess}\begin{pmatrix} A & & B & \\ & C & & D \end{pmatrix}^{(\rho)}$ can be defined as a unique irreducible quotient of

$$\mathrm{Ind}(\delta([\nu_\delta^C\rho, \nu_\delta^D\rho]) \times \delta([\nu_\delta^{C-1}\rho, \nu_\delta^{D-1}\rho]) \times \ldots \times \delta([\nu_\delta^A\rho, \nu_\delta^B\rho])).$$

The content of the paper is as follows. In the second section we recall notation for general linear groups that we use in this paper. The third section recalls some very simple criteria for reducibility or irreducibility of parabolically induced representations. In the fourth section we determine relations between segments defining essentially unitarizable representations $u_{ess}(\delta, n)$. In the fifth section we prove the reducibility criterion in the case when the underlying sets of cuspidal supports are linked, while in the sixth section we prove the criterion when they are not linked. The seventh section gives two formulations of the criterion which we have proved in the previous sections, while the last section clarifies the relation with the work of C. Mœglin and J.-L. Waldspurger.

2. Notation

We recall some notation for general linear groups in the non-archimedean case, following mainly [23], [17] and [30].

2.1. $\mathbb{Z}$-segments in $\mathbb{R}$. By a $\mathbb{Z}$-segment in $\mathbb{R}$, we mean a set of form

$$\{x, x+1, \dots, x+n\},$$

where $x \in \mathbb{R}$ and $n \in \mathbb{Z}_{\geq 0}$. We denote the above set by

$$[x, x+n]_{\mathbb{Z}},$$

or later on, simply by

$$[x, x+n]$$

to shorten notation (this will not cause confusion since we shall not deal with intervals of real numbers in this paper). Then x is called the beginning of Δ, and denoted by $b(\Delta)$, and $x+n$ is called the end of Δ, and denoted by $e(\Delta)$. We denote the set of all $\mathbb{Z}$-segments in $\mathbb{R}$ by $\mathcal{S}(\mathbb{R})$. For $[x,y]_{\mathbb{Z}} \in \mathcal{S}(\mathbb{R})$, let

$$[x,y]_{\mathbb{Z}}^{-} = [x, y-1]_{\mathbb{Z}},$$

$$^{-}[x,y]_{\mathbb{Z}} = [x+1, y]_{\mathbb{Z}}$$

if $x < y$. Otherwise, we take $[x,x]_{\mathbb{Z}}^{-} = {}^{-}[x,x]_{\mathbb{Z}} = \emptyset$.

For $n \in \mathbb{Z}_{>0}$, let

$$\Delta[n] = [-(n-1)/2, (n-1)/2]_{\mathbb{Z}}.$$

Segments $\Delta_1, \Delta_2 \in \mathcal{S}(\mathbb{R})$ are called linked if $\Delta_1 \cup \Delta_2 \in \mathcal{S}(\mathbb{R})$ and $\Delta_1 \cup \Delta_2 \not\in \{\Delta_1, \Delta_2\}$. If the segments Δ_1 and Δ_2 are linked and if Δ_1 and $\Delta_1 \cup \Delta_2$ have the same beginnings, we say that Δ_1 precedes Δ_2, and write

$$\Delta_1 \to \Delta_2.$$

For $\Delta \in \mathcal{S}(\mathbb{R})$ and $x \in \mathbb{R}$, let

$$\Delta_x := \{x + y; y \in \Delta\} \in \mathcal{S}(\mathbb{R}).$$

For a set X, the set of all finite multisets in X is denoted by $M(X)$ (we can view them as all functions $X \to \mathbb{Z}_{\geq 0}$ with finite support; note that finite subsets correspond to all functions $X \to \{0, 1\}$ with finite support). Elements of $M(X)$ are denoted by $(x_1, \dots, x_n)$ (repetitions of elements can occur, and the multiset does not change if we permute x_i's). The number

$$n$$

is called the cardinality of $(x_1, \dots, x_n)$. We call

$$\{x_1, \dots, x_n\}$$

the underlying set of $(x_1, \dots, x_n)$.

The set $M(X)$ has a natural structure of a commutative associative semi group with zero. The operation is denoted additively:

$$(x_1, \dots, x_n) + (y_1, \dots, y_m) = (x_1, \dots, x_n, y_1, \dots, y_m).$$

Take positive integers n and d. Let

$$a(n, d) = (\Delta[d]_{-\frac{n-1}{2}}, \Delta[d]_{-\frac{n-1}{2}+1}, \dots, \Delta[d]_{\frac{n-1}{2}}) \in M(\mathcal{S}(\mathbb{R})). \tag{2.6}$$

2.2. **Groups and representations.** Let F be a non-archimedean locally compact non-discrete field and $|\ \ |_F$ its modulus character. Fix a finite dimensional central division algebra $\mathcal{A}$ over F of rank $d_{\mathcal{A}}$. Denote by $\mathrm{Mat}(n \times n, \mathcal{A})$ the algebra of all $n \times n$ matrices with entries in $\mathcal{A}$. Then $GL(n, \mathcal{A})$ is the group of invertible matrices with the natural topology. The commutator subgroup is denoted by $SL(n, \mathcal{A})$. Denote by

$$\det : GL(n, \mathcal{A}) \to GL(1, \mathcal{A})/SL(1, \mathcal{A})$$

the determinant homomorphism, as defined by J. Dieudonné (for $n = 1$ this is just the quotient map). The kernel is $SL(n, \mathcal{A})$. Denote the reduced norm of $\mathrm{Mat}(n \times n, \mathcal{A})$ by $\mathrm{r.n.}_{\mathrm{Mat}(n\times n,\mathcal{A})/F}$. We identify characters of $GL(n, \mathcal{A})$ with characters of $F^\times$ using $\mathrm{r.n.}_{\mathrm{Mat}(n\times n,\mathcal{A})/F}$. Let

$$\nu = |\mathrm{r.n.}_{\mathrm{Mat}(n\times n,\mathcal{A})/F}|_F : GL(n, \mathcal{A}) \to \mathbb{R}^\times.$$

Denote by

$$G_n$$

the general linear groups $GL(n, \mathcal{A})$ for $n \geq 0$ (we take G_0 to be the trivial group; we consider it formally as the group of 0×0 matrices). The category of all smooth representations of G_n is denoted by $\mathrm{Alg}(G_n)$. The set of all equivalence classes of irreducible smooth representations of G_n is denoted by $\tilde{G}_n$. The subset of unitarizable classes in $\tilde{G}_n$ is denoted by $\hat{G}_n$. The Grothendieck group of the category of all smooth representations of G_n of finite length is denoted by R_n. It is a free $\mathbb{Z}$-module with basis $\tilde{G}_n$. The set of all finite sums in R_n of elements of the basis $\tilde{G}_n$ is denoted by $(R_n)_+$. Set

$$\begin{aligned}
&\mathrm{Irr} = \cup_{n\in\mathbb{Z}_{\geq 0}}\tilde{G}_n,\\
&\mathrm{Irr}^u = \cup_{n\in\mathbb{Z}_{\geq 0}}\hat{G}_n,\\
&R = \oplus_{n\in\mathbb{Z}_{\geq 0}}R_n,\\
&R_+ = \sum_{n\in\mathbb{Z}_{\geq 0}}(R_n)_+.
\end{aligned}$$

The ordering on R is defined by $r_1 \leq r_2 \iff r_2 - r_1 \in R_+$.

The set of cuspidal classes in $\tilde{G}_n$ is denoted by $\mathcal{C}(G_n)$. Denote

$$\begin{aligned}
&\mathcal{C} = \cup_{n\in\mathbb{Z}_{\geq 1}}\mathcal{C}(G_n),\\
&\mathcal{C}^u = \mathcal{C} \cap \mathrm{Irr}^u.
\end{aligned}$$

Let σ_1 and σ_2 be smooth representations of G_{n_1} and G_{n_2}, respectively. Set

$$M_{(n_1,n_2)} := \left\{\begin{bmatrix} g_1 & * \\ 0 & g_2\end{bmatrix}; g_i \in G_i\right\} \subseteq G_{n_1+n_2}.$$

Consider $\sigma_1 \otimes \sigma_2$ as a representation of $M_{(n_1,n_2)}$:

$$\begin{bmatrix} g_1 & * \\ 0 & g_2\end{bmatrix} \mapsto \sigma_1(g_1) \otimes \sigma_2(g_2).$$

Denote by

$$\sigma_1 \times \sigma_2$$

the representation of $G_{n_1+n_2}$ parabolically induced by $\sigma_1 \otimes \sigma_2$ from $M_{(n_1,n_2)}$ (the induction that we consider here is smooth and normalized). Then for three representations, we have

$$(\sigma_1 \times \sigma_2) \times \sigma_3 \cong \sigma_1 \times (\sigma_2 \times \sigma_3).$$

Since the induction functor is exact, we can lift it in a natural way to a $\mathbb{Z}$-bilinear mapping $\times : R_{n_1} \times R_{n_2} \to R_{n_1+n_2}$, and further to $\times : R \times R \to R$. In this way R becomes graded commutative ring.

For $\rho \in \mathcal{C}$ denote by

$$s_\rho$$

the minimal non-negative number such that $\rho \times \nu^{s_\rho}\rho$ reduces. Then $s_\rho \in \mathbb{Z}_{\geq 1}$, and it divides $d_{\mathcal{A}}$ (it can be described in terms of Jacquet-Langlands correspondence established in [11]). Put

$$\nu_\rho := \nu^{s_\rho}.$$

2.3. **Segments in cuspidal representations $\mathcal{C}$.** Let $\Delta \in \mathcal{S}(\mathbb{R})$ and $\rho \in \mathcal{C}$. Set

$$\Delta^{(\rho)} := \{\nu_\rho^x \rho; x \in \Delta\}.$$

The set $\Delta^{(\rho)}$ is called a segment in $\mathcal{C}$. Once we fix ρ, then we call elements in Δ the exponents of elements in $\Delta^{(\rho)}$. Then, when we work with $\Delta^{(\rho)}$, we often drop the superscript (ρ), and instead of $\Delta^{(\rho)}$ and its elements, we refer simply to Δ and its elements.

The set of all segments in $\mathcal{C}$ is denoted by $\mathcal{S}(\mathcal{C})$. We take $\emptyset^{(\rho)} = \emptyset$. For $\Delta^{(\rho)} \in \mathcal{S}(\mathcal{C})$, where $\Delta \in \mathcal{S}(\mathbb{R})$ and $\rho \in \mathcal{C}$, we define

$$(\Delta^{(\rho)})^- := (\Delta^-)^{(\rho)},$$
$$^-(\Delta^{(\rho)}) := (\,^-\Delta)^{(\rho)}.$$

For two segments $\Gamma_1, \Gamma_2 \in \mathcal{S}(\mathcal{C})$, we say that they are linked if there exist linked segments Δ_1, Δ_2 in $\mathcal{S}(\mathbb{R})$ and $\rho \in \mathcal{C}$ such that

$$\Gamma_i = \Delta_i^{(\rho)}, \quad i = 1, 2.$$

In that case we say that Γ_1 precedes Γ_2 if Δ_1 precedes Δ_2, and we then write

$$\Gamma_1 \to \Gamma_2.$$

For $a = (\Delta_1, \dots, \Delta_n) \in M(\mathcal{S}(\mathbb{R}))$ and $\rho \in \mathcal{C}$, set

$$a^{(\rho)} := (\Delta_1^{(\rho)}, \dots, \Delta_n^{(\rho)}) \in M(\mathcal{S}(\mathcal{C})).$$

Let $b = (\Gamma_1, \dots, \Gamma_n) \in M(\mathcal{S}(\mathcal{C}))$ and suppose that Γ_i and Γ_j are linked for some $1 \leq i < j \leq n$. Denote by c the multiset that we get by replacing Γ_i and Γ_j by $\Gamma_i \cup \Gamma_j$ and $\Gamma_i \cap \Gamma_j$ in b (we omit $\emptyset$ if if $\Gamma_i \cap \Gamma_j = \emptyset$). Then we write

$$c \prec b.$$

For $b_1, b_2 \in M(\mathcal{S}(\mathcal{C}))$ we write $b_1 \leq b_2$ if $b_1 = b_2$, or if there exist $c_1, \dots, c_k \in M(\mathcal{S}(\mathcal{C}))$, with $k \geq 2$ such that

$$b_1 = c_1 \prec c_2 \prec \dots \prec c_k = b_2.$$

Then $\leq$ is an ordering on $M(\mathcal{S}(\mathcal{C}))$.

For $\Gamma \in \mathcal{S}(\mathcal{C})$ we define $\operatorname{supp}(\Gamma)$ to be Γ, but considered as an element of $M(\mathcal{C})$. For $a = (\Gamma_1, \dots, \Gamma_n) \in M(\mathcal{S}(\mathcal{C}))$ we define

$$\operatorname{supp}(a) = \sum_{i=0}^{n} \operatorname{supp}(\Gamma_i) \in M(\mathcal{C}).$$

The contragredient representation of π is denoted by $\tilde{\pi}$. For $\Delta \in \mathcal{S}(\mathcal{C})$, set $\tilde{\Delta} := \{\tilde{\rho}; \rho \in \Delta\}$. If $a = (\Delta_1, \dots, \Delta_k) \in M(\mathcal{S}(\mathcal{C}))$, then we put

$$\tilde{a} = (\tilde{\Delta}_1, \dots, \tilde{\Delta}_k).$$

2.4. **Classifications of non-unitary duals.** Let $\Delta = \{\rho, \nu_\rho\rho, \ldots, \nu_\rho^n\rho\} \in \mathcal{S}(\mathcal{C})$. Then the representation

$$\rho \times \nu_\rho^\rho \times \ldots \times \nu_\rho^n\rho$$

has a unique irreducible subrepresentation, which is denoted by

$$\mathfrak{z}(\Delta),$$

and a unique irreducible quotient, which is denoted by

$$\delta(\Delta).$$

Let $a = (\Delta_1, \ldots, \Delta_n) \in M(\mathcal{S}(\mathcal{C}))$. We choose an enumeration of Δ_i's such that for all $i, j \in \{1, 2, \ldots, n\}$ the following holds:

$$\text{if } \Delta_i \to \Delta_j, \text{ then } j < i.$$

Then the representations

$$\begin{aligned}\zeta(a) &:= \mathfrak{z}(\Delta_1) \times \mathfrak{z}(\Delta_2) \times \ldots \times \mathfrak{z}(\Delta_n),\\ \lambda(a) &:= \delta(\Delta_1) \times \delta(\Delta_2) \times \ldots \times \delta(\Delta_n)\end{aligned}$$

are determined by a up to an isomorphism (i.e., their isomorphism classes do not depend on the enumeration). The representation $\zeta(a)$ has a unique irreducible subrepresentation, which is denoted by

$$Z(a),$$

while the representation $\lambda(a)$ has a unique irreducible quotient, which is denoted by

$$L(a).$$

In this way we obtain mappings

$$Z, L : M(\mathcal{S}(\mathcal{C})) \to \mathrm{Irr},$$

which are bijections. Here, Z is called Zelevinsky classification of Irr, while L is called Langlands classification of Irr (we follow the presentation of these classifications given by F. Rodier in [17]). Ror contragredient representations, we

$$L(a)\tilde{} = L(\tilde{a}) \text{ and } Z(a)\tilde{} = Z(\tilde{a}).$$

Denote by $\mathcal{D}$ the set of all essentially square integrable modulo center classes in $\mathrm{Irr}\backslash\hat{G}_0$, and by $\mathcal{D}^u$ the subset of all unitarizable classes in $\mathcal{D}$ (i.e., those having unitary central character). The mapping

$$(\rho, n) \mapsto \delta(\Delta[n]^{(\rho)}), \quad \mathcal{C} \times \mathbb{Z}_{\geq 1} \to \mathcal{D} \tag{2.7}$$

is a bijection.

If $\delta = \delta(\Delta[n]^{(\rho)}) \in \mathcal{D}$, then we denote

$$\nu_\delta = \nu_\rho$$

(we could define ν_δ in the same way as ν_ρ).

For $\delta \in \mathcal{D}$ define $\delta^u \in \mathcal{D}^u$ and $e(\delta) \in \mathbb{R}$ by the following requirement:

$$\delta = \nu_\delta^{e(\delta)} \delta^u.$$

Let $\delta \in M(\mathcal{D})$. We can choose an enumeration of elements of d which satisfies:

$$e(\delta_1) \geq e(\delta_2) \geq \cdots \geq e(\delta_n).$$

Let

$$\lambda(d) = \delta_1 \times \delta_2 \times \ldots \times \delta_n.$$

Then the representation $\lambda(d)$ has a unique irreducible quotient, denoted by $L(d)$. Again $L : M(\mathcal{D}) \to \mathrm{Irr}$ is a bijection, and it is one of the versions of Langlands classification in this case.

The representations

$$u(\delta, n) = L((\nu_\delta^{\frac{n-1}{2}} \delta, \nu_\delta^{\frac{n-1}{2}-1} \delta, \ldots, \nu_\delta^{-\frac{n-1}{2}} \delta)), \quad \delta \in \mathcal{D},$$

are essentially unitarizable (i.e., they become unitarizable after a twist by the appropriate character; see [6] and [4]).

2.5. **Duality - Zelevinsky involution.** Define a mapping

$$^t : \mathrm{Irr} \to \mathrm{Irr}$$

by $t(Z(a)) = L(a), a \in M(\mathcal{S}(\mathcal{C}))$. Extend t additively to R. Clearly, t is a positive mapping, i.e., satisfies: $r_1 \leq r_2 \implies t(r_1) \leq t(r_2)$. A non-trivial fact is that t is also multiplicative, i.e., a ring homomorphism (see [1], [17] and [18]). Further, t is an involution. For $a \in M(\mathcal{S}(\mathcal{C}))$ we define $a^t \in M(\mathcal{S}(\mathcal{C}))$ by the requirement

$$(L(a))^t = L(a^t).$$

We could also use the Zelevinsky classification to define $^t : M(\mathcal{S}(\mathcal{C})) \to M(\mathcal{S}(\mathcal{C}))$, i.e., we would get the same involutive mapping.

One can find more information about the involution in [17].

2.6. **Algorithm of C. Mœglin and J.-L. Waldspurger.** Let $a \in M(\mathcal{S}(\mathbb{R}))$ and $\rho \in \mathcal{S}(\mathcal{C})$. Then there exists $a^t \in M(\mathcal{S}(\mathbb{R}))$, independent of ρ, such that

$$(a^{(\rho)})^t = (a^t)^{(\rho)}.$$

Now we recall the combinatorial algorithm from [14] describing a^t.

Consider segments Δ in a with maximal $e(\Delta)$. Among these segments, choose one with maximal $b(\Delta)$. Denote it by Δ_1, and denote its end by x.

Now consider segments Δ in a which end with $x-1$, and which are linked with Δ_1 (i.e. which precede Δ_1). Among them, if such segments exist, choose one with maximal $b(\Delta)$. Denote it by Δ_2.

One continues this procedure with ends $x-2$, $x-3$, etc., as long as it is possible. The segments considered in this procedure are $\Delta_1, \dots, \Delta_k$ $(k \geq 1)$.

Let

$$\Gamma_1 = [x-k+1, x] \in M(\mathbb{Z}).$$

Let $a^{\leftarrow}$ be the multiset of $M(\mathcal{S}(\mathbb{R}))$ which we get from a by replacing each Δ_i by Δ_i^-, $i = 1, \dots, k$ (we simply omit those Δ_i^- for which $\Delta_i^- = \emptyset$).

If $a^{\leftarrow}$ is non-empty, we now repeat the above procedure with $a^{\leftarrow}$. In this way we get a segment Γ_2 and $(a^{\leftarrow})^{\leftarrow} \in M(\mathcal{S}(\mathbb{R}))$.

Continuing this procedure as long as possible, we get $\Gamma_1, \dots, \Gamma_m \in \mathcal{S}(\mathbb{R})$. Then by [14] (see also [7])

$$a^t = (\Gamma_1, \dots, \Gamma_m).$$

This algorithm will be denoted by

$$\mathrm{MWA}^{\leftarrow}.$$

The set of stages of the algorithm, which end with some segment Γ_i, will be called a step of the algorithm.

2.7. **Dual algorithm.** Lift the contragredient mapping $\tilde{}$ on irreducible representations to a homomorphism $\tilde{} : R \to R$. Then it is a ring homomorphism. One directly sees on generators $\delta(\Delta)$, $\Delta \in \mathcal{S}(\mathcal{C})$, that this homomorphism commutes with the Zelevinsky involution. Thus

$$(L(a)\tilde{})^t = (L(a)^t)\tilde{}, \ a \in M(\mathcal{S}(\mathcal{C})).$$

Therefore, for $a \in M(\mathcal{S}(\mathbb{R}))$ and $\rho \in \mathcal{C}$, we can apply the above algorithm to $L(a^{(\rho)})\tilde{}$, and after that apply once again the contragredient mapping. Since the contragredient mapping is involution, we get the Zelevinsky involution. This gives the following (dual) version of the above algorithm.

Consider segments Δ in a with minimal $b(\Delta)$. Among them, pick a segment with minimal $e(\Delta)$. Denote it by Δ_1, and its beginning with x. Now consider segments Δ in a which begin with $x+1$ and which are linked with Δ_1, if any such segment exists. Among these segments choose one with minimal $e(\Delta)$. Denote it by Δ_2. One continues this procedure with beginnings $x+2$, $x+3$, etc., as long as it is possible. The segments that have shown up in this procedure are denoted by $\Delta_1, \dots, \Delta_k$ $(k \geq 1)$. Put

$$\Gamma_1 = [x, x+k-1] \in M(\mathbb{Z}).$$

Let ${}^{\rightarrow}a$ be the multiset of $M(\mathcal{S}(\mathbb{R}))$ which we get from a by replacing each Δ_i by ${}^-\Delta_i$ (if ${}^-\Delta_i = \emptyset$, we simply omit it). If ${}^{\rightarrow}a$ is non-empty, we repeat the above procedure with ${}^{\rightarrow}a$.

In this way we get Γ_2 and ${}^{\to}({}^{\to}a)$. Continuing this procedure as long as possible, we get $\Gamma_1, \dots, \Gamma_m \in \mathcal{S}(\mathbb{R})$. Then

$$a^t = (\Gamma_1, \dots, \Gamma_m).$$

This algorithm will be denoted by

$${}^{\to}\mathrm{MWA}.$$

We shall usually apply the above algorithm(s) to elements of $M(\mathcal{S}(\mathcal{C}))$ in an obvious way. It is easy to show that

$$(a(n,d)^{(\rho)})^t = a(d,n)^{(\rho)}.$$

2.8. **Upper bound for the lengths of the segments in the dual multisegment.** We later use the following observation of C. Mœglin and J.-L. Waldspurger from [14] (see also [3]): if there exists a segment Δ of length m such that all the ends of segments in $a \in M(\mathcal{S}(\mathcal{C}))$ are contained in Δ, then the length of segments in a^t can be at most m.

Dually, we get the following observation: If there is a segment Δ of length m such that all the beginnings of segments in $a \in M(\mathcal{S}(\mathcal{C}))$ are contained in Δ, then the length of segments in a^t is at most m.

2.9. **Support.** Let $\pi \in \mathrm{Irr}$. Take $a \in M(\mathcal{S}(\mathcal{C}))$ such that $\pi = L(a)$. Then the support of π is defined by

$$\mathrm{supp}(\pi) = \mathrm{supp}(a).$$

If we take $b \in M(\mathcal{S}(\mathcal{C}))$ such that $\pi = Z(a)$, then $\mathrm{supp}(\pi) = \mathrm{supp}(b)$.

Suppose that for a finite length representation π', for each two irreducible subquotients π_1 and π_2 one has $\mathrm{supp}(\pi_1) = \mathrm{supp}(\pi_2)$. Then we define $\mathrm{supp}(\pi')$ to be $\mathrm{supp}(\pi)$, where π is (any) irreducible subquotient of π'.

2.10. **Classification or the unitary dual.** Denote by

$$B_{\mathrm{rigid}} = \{u(\delta, n); \delta \in \mathcal{D}^u, n \in \mathbb{Z}_{\geq 1}\}.$$

and

$$B = B_{\mathrm{rigid}} \cup \{\nu_\delta^{\alpha} u(\delta, n) \times \nu_\delta^{-\alpha} u(\delta, n); u(\delta, n) \in B_{\mathrm{rigid}}, 0 < \alpha < 1/2\}.$$

Then the unitary dual is described by the following:

Theorem 2.1. *(* [4],[6],[19],[23]*)*

(1) *Let $\tau_1, \dots, \tau_n \in B$. Then the representation*

$$\pi := \tau_1 \times \dots \times \tau_n$$

is irreducible and unitary.

(2) *Suppose that a representation* π' *is obtained from* $\tau'_1, \dots, \tau'_{n'} \in B$ *in the same manner as* π *was obtained from* $\tau_1, \dots, \tau_n$ *in (1). Then* $\pi \cong \pi'$ *if and only if* $n = n'$ *and the sequences* $(\tau_1, \dots, \tau_n)$ *and* $(\tau'_1, \dots, \tau'_n)$ *coincide after a renumeration.*

(3) *Each irreducible unitary representation of* $GL(m, F)$, *for any* m, *can be obtained as in (1).*

Recall that

$$u(\Delta[d]^{(\rho)}, n) = L(a(d, n)^{(\rho)}),$$

and

$$Z(a(n, d)^{(\rho)}) \cong L(a(d, n)^{(\rho)}) \tag{2.8}$$

for $n, d \in \mathbb{Z}_{\geq 0}$ and $\rho \in \mathcal{C}$.

3. Some criteria for reducibility and irreducibility

3.1. **A reducibility criterion.** Let $a, b \in M(\mathcal{D})$. We know

$$L(a + b) \leq L(a) \times L(b).$$

Thus

$$L((a + b)^t) = L(a + b)^t \leq L(a)^t \times L(b)^t.$$

Also

$$L(a^t + b^t) \leq L(a^t) \times L(b^t) = L(a)^t \times L(b)^t.$$

This implies the following well known reducibility criterion

$$\text{If } (a + b)^t \neq a^t + b^t, \text{ then } L(a) \times L(b) \text{ reduces.} \tag{3.9}$$

In general, $(a + b)^t = a^t + b^t$ does not imply the irreducibility of $L(a) \times L(b)$.

3.2. **Irreducibility criterion of I. Badulescu.** Suppose that $a_1, a_2 \in M(\mathcal{S}(\mathcal{C}))$ satisfy $(a_1 + a_2)^t = a_1^t + a^t$ (if this is not the case, then $L(a_1) \times L(a_2)$ is reducible). Assume that for each $b \in M(\mathcal{S}(\mathcal{C}))$ holds

$$b < a_1 + a_2 \implies b^t \not< (a_1 + a_2)^t. \tag{3.10}$$

Then $L(a_1) \times L(a_2)$ is irreducible.

In general, irreducibility of $L(a_1) \times L(a_2)$ does not imply that (3.10) holds.

For the convenience of the reader, we repeat the argument from [3].

Suppose that $L(a_1) \times L(a_2)$ is reducible. Then in R we have

$$L(a_1) \times L(a_2) = L(a_1 + a_2) + \sum_{i=1}^{k} n_i L(b_i),$$

where $n_i \in \mathbb{Z}_{\geq 1}$, $k \geq 1$ and

$$b_i < a_1 + a_2, i = 1, \ldots, k.$$

Applying the Zelevinsky involution, we get (in R)

$$L(a_1^t) \times L(a_2^t) = L(a_1^t + a_2^t) + \sum_{i=1}^{k} n_i L(b_i)^t$$

(here we have used $(a_1 + a_2)^t = a_1^t + a^t$). The properties of the Langlands classification imply

$$b_1^t, \ldots, b_k^t < a_1^t + a_2^t.$$

This contradicts the assumptions of the criterion.

3.3. **Contacting and crossing.** To present the next criterion, we need the following:

Definition 3.1. *Let $a, b \in M(\mathcal{D})$. We say that a is in contact with b (or simply that a contacts b), if there exist segments Δ_1 and Δ_2 in a and b respectively, which are juxtaposed (two non-empty segments are called juxtaposed if they are disjoint and if their union is a segment).*

We say that a and b are crossed if a contacts b^t, and a^t contacts b.

We say that irreducible representations are in contact (resp., are crossed) if the segments corresponding to them with respect to the Langlands classification are in contact (resp., are crossed).

Remark 3.2. *Observe that a is in contact with b if and only if $\tilde{a}$ in contacts with $\tilde{b}$. Further, the action of Zelevinsky involution and the contragredient mapping commute on segments, i.e., $(\tilde{\Delta})^t = (\Delta^t)\tilde{}$, which implies that they commute on R. From this, it follows easily that*

$$(\tilde{a})^t = (a^t)\tilde{}.$$

Therefore,

$$a \text{ and } b \text{ are crossed if and only if } \tilde{a} \text{ and } \tilde{b} \text{ are crossed.}$$

Clearly $L(a) \times L(b)$ reduces $\iff$ $L(\tilde{a}) \times L(\tilde{b})$ reduces $\iff$ $L(a^t) \times L(b^t)$ reduces $\iff$ $L(\tilde{a}^t) \times L(\tilde{b}^t)$ reduces.

These equivalences also hold for the Zelevinsky classification (and $L(a) \times L(b)$ reduces $\iff$ $Z(a) \times Z(b)$ reduces).

3.4. Irreducibility criterion of I. Badulescu, E. Lapid and A. Míngues.

This criterion for $a, b \in M(\mathcal{D})$ says the following:

$$\text{If } a \text{ is not in contact with } b\text{, then } L(a) \times L(b)^t \text{ is irreducible.} \tag{3.11}$$

We get directly from this the following criterion:

$$\text{If } a \text{ and } b^t\text{, or } a^t \text{ and } b\text{, are not in contact, then } L(a) \times L(b) \text{ is irreducible.} \tag{3.12}$$

In other words:

$$\text{If } L(a) \times L(b) \text{ is reducible, then } a \text{ and } b \text{ are crossed.} \tag{3.13}$$

4. CONTACTS OF NON-INDUCED ESSENTIALLY UNITARIZABLE REPRESENTATIONS

We now describe when two representations $L(a(n,d)^{(\nu^k\rho)}$ and $L(a(m,d)^{(\nu^l\rho;)})$, supported on the same cuspidal line, are in contact.

4.1. Contacts among $L(a(n,d)^{(\rho)})$'s.

Let

$$\pi_1 = L(a(n,d)^{(\nu_{\rho'}^k\rho')}), \qquad \pi_2 = L(a(m,e)^{(\nu_{\rho'}^l\rho')}).$$

After twisting ρ', we can write

$$L(a(n,d)^{(\nu_{\rho'}^k\rho')}) = L([1,d]^{(\rho)}, [2,d+1]^{(\rho)}], \dots, [n, d+n-1]^{(\rho)}),$$

$$L(a(m,e)^{(\nu_{\rho'}^l\rho')}) = L([b, b+e-1]^{(\rho)}, [b+1, b+e]^{(\rho)}, \dots, [b+m-1, b+m-1+e-1]^{(\rho)}),$$

for some $\rho \in \mathcal{C}$.

In what follows we assume

$$b \in \mathbb{Z}$$

(otherwise, $\pi_1 \times \pi_2$ is always irreducible).

Now π_1 and π_2 are in contact if and only if

$$[1,n] \cap [b+e, b+m-1+e] \neq \emptyset \quad \text{or} \quad [d+1, d+n] \cap [b, b+m-1] \neq \emptyset.$$

We can graphically interpret this by the following drawing:





Looking at the above drawing, we have contact between π_1 and π_2 if and only if the intersection of (projections to the horizontal axis of) bold lines is non-empty, or the intersection of (projections of) dashed lines is non-empty.

We study when $\pi_1 \times \pi_2$ reduces. Since $\pi_1 \times \pi_2$ reduces if and only if $\pi_2 \times \pi_1$ reduces, without lost of generality we can always enumerate π_i's in such a way that

$$1 \leq b.$$

Since we are interested in reducibility of $\pi_1 \times \pi_2$, we consider only the case when the union of underlying sets of supports of π_1 and π_2 is a segment (if it is not, then $\pi_1 \times \pi_2$ is irreducible). Therefore, we assume

$$b \leq d+n.$$

We retain the assumptions

$$1 \leq b \leq d+n \tag{4.14}$$

in what follows.

Observe that the first condition for π_i's to be in contact, $[1,n] \cap [b+e, b+m-1+e] \neq \emptyset$, is now equivalent to $b+e \leq n$, i.e.,

$$b+e-1 < n.$$

4.2. **Another notation.** We denote
$A_1 = 1, \qquad B_1 = d,$

$$C_1 = n, \qquad D_1 = n+d-1,$$

$$A_2 = b, \qquad B_2 = b+e-1,$$

$$C_2 = b+m-1, \qquad D_2 = b+m-1+e-1.$$

Obviously, $A_1 \leq B_1, C_1 \leq D_1$, $A_2 \leq B_2, C_2 \leq D_2$, and

$$B_1 - A_1 = D_1 - C_1,$$

$$B_2 - A_2 = D_2 - C_2.$$

The previous assumption, $1 \leq b \leq d+n$, now becomes

$$A_1 \leq A_2 \leq D_1 + 1. \tag{4.15}$$

Now we have contact if and only if

$$[A_1, C_1] \cap [B_2+1, D_2+1] \neq \emptyset \quad \text{or} \quad [B_1+1, D_1+1] \cap [A_2, C_2] \neq \emptyset.$$

The previous drawing now corresponds to the following drawing in the new notation:



We have contact if and only if the (projections to the horizontal axis of) bold lines have non-empty intersection, or the dashed lines have non-empty intersection.

Since we assume $A_1 \leq A_2$, the first condition is equivalent to $B_2 + 1 \leq C_1$, i.e., $B_2 < C_1$. Therefore, we have contact if and only if

$$B_2 < C_1 \quad \text{or} \quad [B_1+1, D_1+1] \cap [A_2, C_2] \neq \emptyset. \tag{4.16}$$

Remark 4.1. *Let the underlying sets of supp*$(L(a(n,d)^{(\nu^k_{\rho'},\rho')})$ *and supp*$(L(a(m,e)^{(\nu^l_{\rho'},\rho')})$ *be linked segments. This implies*

$$A_1 < A_2, D_1 < D_2, A_2 \leq D_1 + 1.$$

Suppose that the π_i*'s are in contact, and that the intersection of bold segments is non-empty. This implies*

$$B_2 + 1 \leq C_1 \text{ i.e. } B_2 < C_1.$$

We then show that

$$[B_1+1, D_1+1] \cap [A_2, C_2] \tag{4.17}$$

is also non-empty (which is the intersection of the dashed segments). We consider two cases.

1. *Suppose* $B_1 + 1 \leq A_2$. *Then* (4.17) *is non-empty if and only if* $[A_2, D_1 + 1] \neq \emptyset$. *Clearly, this is the case if and only if* $A_2 \leq D_1 + 1$. *We know that this holds. Therefore,* (4.17) *is non-empty.*
2. *Now, suppose* $A_2 < B_1 + 1$. *Then* (4.17) *is non-empty if and only if* $[B_1 + 1, C_2] \neq \emptyset$, *which is the case if and only if* $B_1 + 1 \leq C_2$. *We show below that this holds.*

   Observe that $A_1 < A_2$, $D_1 < D_2$ *and* $B_2 + 1 \leq C_1$ *imply* $A_1 + B_2 + D_1 + 1 < A_2 + C_1 + D_2$, *i.e.,* $A_1 - C_1 + D_1 + 1 < A_2 - B_2 + D_2$. *From* $A_i + D_i = B_i + C_i$, $i = 1, 2$, *we get* $B_1 + 1 < C_2$. *This obviously implies the inequality* $\leq$, *which we wanted to prove. Therefore,* (4.17) *is not empty.*

The proof of the claim of the remark is now complete. In this way we have proved that two multisegments as above are in contact if and only if the dashed segments intersect.

5. Linked case

We continue with the notation introduced in the previous section.

Proposition 5.1. *Suppose that the underlying sets of*

$$supp(L(a(n,d)^{(\nu^k_{\rho'}\rho')}) \quad and \quad supp(L(a(m,e)^{(\nu^l_{\rho'}\rho')})$$

are linked segments. Then

$$L(a(n,d)^{(\nu^k_{\rho'}\rho')}) \times L(a(m,e)^{(\nu^l_{\rho'}\rho')}) \tag{5.18}$$

reduces if and only if $L(a(n,d)^{(\nu^k_{\rho'}\rho')})$ *and* $L(a(m,e)^{(\nu^l_{\rho'}\rho')})$ *are crossed.*

Proof. Thanks to (3.13), we know that the reducibility of (5.18) implies that the corresponding multisegments are crossed. We need to prove the opposite implication, i.e. that if we have crossed multisegments in the lemma, then we have reducibility.

Let

$$\pi_1 = L(a(n,d)^{(\nu^k_{\rho'}\rho')}), \qquad \pi_2 = L(a(m,e)^{(\nu^l_{\rho'}\rho')}),$$
$$a_1 = a(n,d)^{(\nu^k_{\rho'}\rho')}, \qquad a_2 = a(m,e)^{(\nu^l_{\rho'}\rho')}.$$

Here we can write

$$L(a(n,d)^{(\nu^k_{\rho'}\rho')} = L([1,d]^{(\rho)}, [2,d+1]^{(\rho)}], \dots, [n, d+n-1]^{(\rho)}]),$$

$$L(a(m,e)^{(\nu^l_{\rho'}\rho')}) = L([b, b+e-1]^{(\rho)}, [b+1, b+e]^{(\rho)}, \dots, [b+m-1, b+m-1+e-1]^{(\rho)}]),$$

for some $\rho \in \mathcal{C}$, where $1 < b$, $d + n < b + m - 1 + e$, and $b \leq d + n$.

We introduce A_i, B_i, C_i, D_i by the same formulas as in 4.2. The linking condition in this notation is

$$A_1 < A_2,$$
$$D_1 < D_2,$$
$$A_2 \leq D_1 + 1.$$

Further, by Remark 4.1, the crossing condition is equivalent to

$$[B_1 + 1, D_1 + 1] \cap [A_2, B_2] \neq \emptyset$$

and

$$[C_1 + 1, D_1 + 1] \cap [A_2, C_2] \neq \emptyset.$$

Since $A_2 \leq D_1 + 1$, the above two conditions are equivalent to

$$B_1 + 1 \leq B_2$$

and

$$C_1 + 1 \leq C_2.$$

We consider several cases.

(1) Let $D_1 < B_2$.
   (a) Suppose $C_1 < A_2$.
   We first illustrate the situation graphically:



(5.19)



(5.20)



(5.21)

We now show $a_1^t + a_2^t \neq (a_1+a_2)^t$. Then the reducibility criterion (3.9) implies reducibility. Observe that $a_1^t + a_2^t = a(d,n)^{(\nu^k_{\rho'}\rho')} + a(e,m)^{(\nu^l_{\rho'}\rho')}$. Therefore in this multisegment, there are segments of length n and m. The assumptions $C_1 < A_2$ and $D_1 < B_2$, together with MWA$^{\leftarrow}$ directly imply that $(a_1+a_2)^t$ will have at least one segment of length $n+m$ (see the graphical interpretation). Namely, in the first $B_2 - D_1 - 1$ steps of the algorithm, we get segments of length m, and in the following step we get a segment of length $m+n$. This completes the proof of reducibility in this case.

(b) Now suppose $A_2 \leq C_1$.
We again illustrate the situation graphically:



(5.22)



(5.23)

Observe that in $a_1^t + a_2^t$ there is only one segment starting at exponent $A_1 = 1$. This is the segment $[A_1, C_1]^{(\rho)} = [1,n]^{(\rho)}$.
Now applying $^{\rightarrow}$MWA, starting with exponent 1, we get exponents 2, 3, …, n. Since $n = C_1 < C_2$, we can find a segment in a_2 starting with $n+1$. The assumption $D_1 < B_2$ implies that this segment is linked with the previous segment used in the algorithm. This implies that the (unique) segment in $(a_1+a_2)^t$ starting with A_1 is not $[A_1, C_1]^{(\rho)} = [1,n]^{(\rho)}$, as was the the case in $a_1^t + a_2^t$. Thus $(a_1+a_2)^t \neq a_1^t + a_2^t$, which implies reducibility.

(2) Let $B_2 \leq D_1$.

(a) Suppose $C_1 < A_2$.
We again illustrate the situation graphically:



(5.24)

In $a_1^t + a_2^t$ there is a unique segment which ends with exponent D_2. It is $[B_2, D_2]^{(\rho)}$.

We now start MWA$^{\leftarrow}$. It starts with exponent D_2, and proceeds with $D_2 - 1, \ldots, B_2$. Since $B_1 < B_2$, we can find a segment in a_2 ending with $B_2 - 1$. Since $C_1 < A_2$, this segments precedes the previous one ending with B_2. This implies that the segment in $(a_1 + a_2)^t$ ending with exponent D_2 is not $[B_2, D_2]^{(\rho)}$. This implies $a_1^t + a_2^t \neq (a_1 + a_2)^t$, which again implies reducibility.

(b) Suppose now $A_2 \leq C_1$.

We again illustrate the situation graphically:



(5.25)

If $e \leq d$, then start MWA$^{\leftarrow}$, and the first segment that we get from the algorithm implies $a_1^t + a_2^t \neq (a_1 + a_2)^t$, which implies reducibility.

If $d \leq e$, then start $^{\rightarrow}$MWA, and the first segment that we get from the algorithm implies $a_1^t + a_2^t \neq (a_1 + a_2)^t$. Again we get reducibility.

$\square$

6. Non-linked case

We continue with the notation of the last two sections.

In this section, we consider the case where the underlying set of $\mathrm{supp}(L(a(n,d)^{(\nu^k_{\rho'},\rho')})$ contains the underlying set of $\mathrm{supp}(L(a(m,e)^{(\nu^l_{\rho'},\rho')})$. Our aim in this section is to prove that

$$L(a(n,d)^{(\nu^k_{\rho'},\rho')}) \times L(a(m,e)^{(\nu^l_{\rho'},\rho')}) \tag{6.26}$$

is irreducible. We know by criterion (3.11) that (6.26) is irreducible if $L(a(n,d)^{(\nu^k_{\rho'},\rho')})$ and $L(a(m,e)^{(\nu^l_{\rho'},\rho')})$ are not crossed.

This is the reason that we assume that these two representations are crossed in what follows.

Since $L(a(n,d)^{(\nu^k_{\rho'},\rho')}) \times L(a(m,e)^{(\nu^l_{\rho'},\rho')})$ is irreducible if and only the dual representation $L(a(d,n)^{(\nu^k_{\rho'},\rho')}) \times L(a(e,m)^{(\nu^l_{\rho'},\rho')})$ is irreducible, it is enough to consider the case

$$n \leq d.$$

Below we use the notation π_i, a_i, A_i, B_i, C_i, D_i, $i = 1, 2$, from the previous section.

Now the condition of inclusion of underlying sets tells us in this notation,

$$A_1 \leq A_2,$$

$$D_2 \leq D_1,$$

while the assumption $n \leq d$ becomes

$$C_1 \leq B_1.$$

Further, the segments are crossed if and only if both conditions below hold:

$$C_2 + 1 \leq C_1 \text{ or } [B_1 + 1, D_1 + 1] \cap [A_2, B_2] \neq \emptyset; \tag{1}$$

$$B_2 + 1 \leq B_1 \text{ or } [C_1 + 1, D_1 + 1] \cap [A_2, C_2] \neq \emptyset. \tag{2}$$

Since we assume that a_1 and a_2 are crossed, we assume that both above conditions hold in what follows. Since $B_2, C_2 < D_1 + 1$, the above requirements are equivalent to

$$C_2 + 1 \leq C_1 \text{ or } B_1 + 1 \leq B_2 \text{ (i.e., } C_2 < C_1 \text{ or } B_1 < B_2\text{)}; \tag{1}$$

$$B_2 + 1 \leq B_1 \text{ or } C_1 + 1 \leq C_2 \text{ (i.e., } B_2 < B_1 \text{ or } C_1 < C_2\text{)}. \tag{2}$$

Thus the crossing condition is

$$C_2 < C_1 \text{ and } B_2 < B_1 \quad \text{or} \quad C_1 < C_2 \text{ and } B_1 < B_2.$$

Without lost of generality, we can assume

$$A_2 + D_2 \leq A_1 + D_1$$

(if this is not the case, passing to contragredients will bring us to this case). This implies

$$B_2 + C_2 \leq B_1 + C_1.$$

Therefore, the crossing condition for the case $A_2 + D_2 \leq A_1 + D_1$ is

$$C_2 < C_1 \text{ and } B_2 < B_1.$$

Observe that $B_2 < B_1$ (i.e. $b + e - 1 < d$) implies

$$e < d.$$

Lemma 6.1. *With the above notation and assumptions, we have*

$$a_1^t + a_2^t = (a_1 + a_2)^t.$$

Proof. We treat two cases separately. The first case is

$$D_2 < B_1$$

(i.e. $b + m - 1 + e - 1 < d$).

We illustrate the situation by the drawing below:



(6.27)

Starting $\mathrm{MWA}^{\leftarrow}$ with D_1, after the first step, we get the segment $[B_1, D_1]^{(\rho)}$ of $(a_1 + a_2)^t$. Note that at each stage or first step of $\mathrm{MWA}^{\leftarrow}$, there is only one possibility for the segment that we can use in $\mathrm{MWA}^{\leftarrow}$ (the segments that we are using are $[C_1, D_1]$, …, $[A_1, B_1]$). We continue in the same way - where there is only one possibility for the choice of segment at each stage - as long as possible. We get a sequence of segments $[B_1, D_1]^{(\rho)}$, $[B_1 - 1, D_1 - 1]^{(\rho)}$, …, $[D_2 + 1, D_2 + n]^{(\rho)}$ (recall $D_2 < B_1$).

Now we perform the following step, starting with $D_2 + n - 1$. We get the following $n - 1$ points: $D_2 + n - 1$, $D_2 + n - 2$, …, $D_2 + 1$ (at each stage, there is only one choice of the segment for $\mathrm{MWA}^{\leftarrow}$). Now D_2 is the end of $[1, D_2]$ and $[C_2, D_2]$.

If $1 < C_2$, then $[C_2, D_2]$ does not precede $[2, D_2 + 1]$, while $[1, D_2]$ does. Now, $\mathrm{MWA}^{\leftarrow}$ tells us that we take out the end of $[1, D_2]$. The segment that we get in this step is $[D_2, D_2 + n - 1]^{(\rho)}$. If $1 = C_2$, then $[C_2, D_2] = [1, D_2]$ and we take out the end of the segment $[1, D_2]$ from the part corresponding to a_1. The segment that we get in this step is again $[D_2, D_2 + n - 1]^{(\rho)}$.

We continue this way up to the starting point $C_1 + e - 1$ for the algorithm (the corresponding segment must come from a_1 since $C_2 < C_1$ and now the segments are of the same length). In the previous steps of the algorithm, when we had two segments with the same end, then only one was allowed by $\mathrm{MWA}^{\leftarrow}$, and we took that one (it came from a_1). For $C_1 + e - 1$, we can have two segments, but they are the same, and we choose the one coming from a_1. We continue in this way, and after this step we get the segment $[C_1 + e - n, C_1 + e - 1]^{(\rho)}$ of $(a_1 + a_2)^t$.

We can illustrate the following step of the algorithm by the drawing


(6.28)

We perform the following step, starting with C_1+e-2. It may happen that two segments end with this point. If two of them end with it, then by MWA$^{\leftarrow}$ we need to take the one coming from a_2, since it is shorter. We get the following n points: C_1+e-2, C_1+e-3, $\dots$, $C_1+e-n-1$. At each stage, there is either only one choice for next segment, or there are two choices (both are preceding the previous segment used in the algorithm). Then by MWA$^{\leftarrow}$ we choose the shorter segment, which is coming from a_1.

Further, starting with the points C_1+e-3, $\dots$, C_1, in all the stages of the algorithm, when we had two choices, we chose the end of the segment coming from a_1 because of the order that is considered on segments. In this way, we got the multisegment

$$a(d,n)^{(\nu^k_{\rho'},\rho')} = a_1^t.$$

Continuing further, we obviously get

$$a(d,n)^{(\nu^k_{\rho'},\rho')} + a(e,m)^{(\nu^l_{\rho'},\rho')} = a_1^t + a_2^t.$$

Therefore, in the case $D_2 < B_1$ we have shown that $(a_1+a_2)^t = a_1^t + a_2^t$.

Now we shall consider the case

$$B_1 \le D_2.$$

Recall that we also assume

$$C_2 < C_1, B_2 < B_1 \text{ and } C_1 \le B_1.$$

We illustrate this situation with the following drawing:



(6.29)

We start first step of MWA$^{\leftarrow}$ with D_1 (observe that $B_2 < B_1$ and $C_2 < C_1$ imply $B_2+C_2 < B_1 + C_1$, i.e., $A_2 + D_2 < A_1 + D_1$, which implies $D_2 < D_1$ since $A_1 \leq A_2$). We have only one choice for picking the segment ending with points $D_1 - 1, D_1 - 2, \dots$ until $D_2 + 1$ (including it). At D_2, the preceding condition from MWA$^{\leftarrow}$ tells us that we need to take the end of the segment from a_1 since $e < d$. We perform the same procedure with other the point from the first step.

The procedure is the same as long as segments coming from a_1 are longer than the segments coming from a_2. Further, if the lengths are the same, we have two of the same segments with this end, and we pick the one coming from a_1. Since $C_2 < C_1$, the segment with which we start in this case always comes from a_1. This stops to being the case if the starting point of MWA$^{\leftarrow}$ is $C_1 + e - 2$.

Consider the next step, for the starting point $C_1 + e - 2$. Then the segment coming from a_1 has the length shorter by 1 than the segment coming from a_2.

Suppose $D_2 < C_1 + e - 2$. Then we start with the segment coming from a_1. Now the fact that lengths of segments coming from a_1 are shorter implies that we always pick the end of a segment coming from a_1. We get the segment $[C_1 + e - 1 - n, C_1 + e - 2]^{(\rho)}$ in this way.

Suppose $D_2 = C_1 + e - 2$. Then by MWA$^{\leftarrow}$ we need to start with the segment coming from a_1, and in each further stage, again MWA$^{\leftarrow}$ (the "preceding condition") tells us that we must pick a segment coming from a_1. In this way we get the segment $[D_2 - m + 1, D_2]^{(\rho)}$.

We continue this way, the steps beginning with ends $C_1 + e - i$, $i < 2$, as long as $D_2 < C_1 + e - i$.

For the point D_2, we have two segments ending with D_2. By MWA$^{\leftarrow}$, we need to start with the one coming from a_1, and the procedure continues with segments coming from a_1.

Up to now, MWA$^{\leftarrow}$ has been performed only inside the segments coming from a_1.

In the next step of MWA$^{\leftarrow}$, we again start with D_2. Now the only segment ending with D_2 comes from a_2. Since the segments coming from a_1 are shorter than the ones coming from a_2, the preceding condition of MWA$^{\leftarrow}$ tells us that we must always use segments coming from a_2.

In the next step, we use segments coming from a_1. After this step, the segments coming from a_1 are shorter than the ones coming from a_2. In the following step we must start with the segments coming from a_2.

We continue the steps starting with ends X in this way, as long as $C_1 \leq X$. After this, the remaining segments all come from a_2, and we continue and finish MWA$^{\leftarrow}$ with them.

Observe that each step of MWA$^{\leftarrow}$ was performed completely inside the segments coming from a_1 or a_2. Therefore, we get that $(a_1 + a_2)^t$ is

$$a_1^t + a_2^t.$$

This completes the proof of the lemma. □

Remark 6.2. *Observe that the lemma holds whenever the underlying set of* a_1 *contains the underlying set of* a_2 *(this follows from the fact that* ^{t} *is an involution, the fact that the lemma holds when the corresponding induced representation is irreducible and* (3.11)*).*

Lemma 6.3. *If* $c < a_2$ *then* $c^t \not< a_2^t$.

Proof. Suppose that there exists $c \in M(\mathcal{S}(\mathcal{C}))$ such that $c < a_2$ and $c^t < a_2^t$.

All the segments in a_2^t have length m, and any linking among segments of a_2^t will produce a segment longer than m. Further, linkings may only increase the maximal length of segments. Since $c^t < a_2^t$, we have a segment in c of length at least $m + 1$.

The ends of the segments from a_2 form the segment $[B_2, D_2]^{(\rho)}$. Therefore, the ends of segments from c are contained in $[B_2, D_2]$. By 2.8, this implies that the segments in c^t are not longer than $D_2 - B_2 + 1 = C_2 - A_2 + 1 = m$.

This contradiction completes the proof of the lemma. □

Proposition 6.4. *Suppose that underlying set of supp*$(L(a(n,d)^{(\nu^k_{\rho'},\rho')})$ *contains the underlying set of supp*$(L(a(m,e)^{(\nu^l_{\rho'},\rho')})$*. Then* $L(a(n,d)^{(\nu^k_{\rho'},\rho')}) \times L(a(m,e)^{(\nu^l_{\rho'},\rho')})$ *is irreducible.*

Proof. We continue with the previous notation and the previous assumptions:

$$B_2 < B_1, C_2 < C_1, C_1 \leq B_1 \text{ (i.e., } n \leq d).$$

We suppose that

$$L(a_1) \times L(a_2)$$

reduces. Then there exists

$$f < a_1 + a_2 \tag{6.30}$$

such that $L(f)$ is a subquotient of $L(a_1) \times L(a_2)$. Therefore

$$L(f^t) \text{ is a subquotient of } \ L(a_1^t) \times L(a_2^t).$$

Thus, $f^t \leq a_1^t + a_2^t$. Observe that $f^t = a_1^t + a_2^t$ would contradict the previous lemma since $f \neq a_1 + a_2$. Thus

$$f^t < a_1^t + a_2^t. \tag{6.31}$$

In the rest of the proof we fix some $f < a_1 + a_2$ satisfying $f^t < a_1^t + a_2^t$.

Since $C_2 < C_1$, the beginnings of all segments in $a_1 + a_2$ are contained in $[A_1, C_1]^{(\rho)}$. This also holds for f since $f < a_1 + a_2$ (and since the beginnings of all segments in f are contained in the beginnings of all segments in $a_1 + a_2$). This implies that lengths of segments in f^t are at most n. Since $f^t < a_1^t + a_2^t$, we conclude that no segment from a_1^t can take part in any linking which produces f^t. Therefore

$$f^t = a_1^t + c_t$$

for some $c_t \in M(\mathcal{S}(\mathcal{C}))$. Observe that above considerations imply

$$c_t < a_2^t.$$

We now apply the above arguments to f^t instead of f, and use the fact that $B_2 < B_1$. In the same way we get for $f = (f^t)^t$ that

$$f = a_1 + c$$

for some

$$c < a_2.$$

Now we prove that

$$c_t = c^t.$$

We split the proof of this fact into two parts.

(1) First we consider the case

$$D_2 < C_1,$$

and apply MWA$^{\leftarrow}$ to this case.

We illustrate f and c in this situation by the following example:



(6.32)

We start MWA$^{\leftarrow}$ as before (with D_1), and perform it until at each step there is only one segment which satisfies the algorithm. In this way, we get the following segments: $[C_1, D_1]^{(\rho)}, \dots, [D_2+1, D_2+n]^{(\rho)}$.

In the following step, starting with D_2+n-1, we again get the following $n-1$ points: D_2+n-1, $D_2+n-2, \dots, D_2+1$ (at each stage, there is only one segment which satisfies the algorithm). Now D_2 is the end of $[1, D_2]$, and of $[X, D_2]$ for some X which satisfies $A_2 \leq X \leq C_2$ ($[X, D_2]$ is the segment from c with biggest end). If $1 < X$, then $[X, D_2]$ does not precede $[2, D_2+1]$, while $[1, D_2+1]$ does. We take out the end of $[1, D_2]$ and get $[D_2, D_2+n-1]^{(\rho)}$. If $1 = X$, then $[X, D_2] = [1, D_2+1]$, and we take out the end of the segment $[1, D_2]^{(\rho)}$, considered as a segment from a_1, and again get $[D_2, D_2+n-1]^{(\rho)}$.

We claim that we can perform the first n steps of the algorithm using only segments coming from a_1 (this does not mean that a copy of a segment that we are using is not also in c). Suppose that we have chosen some starting point Y satisfying $n = C_1 \leq Y \leq D_2 - n - 2$, and that in the previous steps we have not used segments from c. Since $D_2 < n = C_1$, from $f^t = a_1^t + c_t$, with $c_t < a_2^t$, we see that this step of MWA$^{\leftarrow}$ must end with $Y - n + 1$. We also know that for the following steps, until the n-th (including), we must get segments of length n.

Suppose that at some stage of the algorithm, starting from Y, we have two possibilities (it follows easily from our inductive assumption that there cannot be more than two, and that one of them must come from a segment coming from a_1 and the other one from a segment from c). Look at the segments whose ends we consider. Now, we have three possibilities. If the segment coming from a_1 is shorter than the one from c, we take the end of the segment coming from a_1 (the lexicographic requirement and "preceding" requirement from MWA$^{\leftarrow}$ are fulfilled). If the segments are of the same length, they are equal, and we take the end of the one coming from a_1. The last possibility is that the segment coming from a_1 is longer then the one from c. But then the "preceding" requirement tells us that we need to take the end of the segment coming from a_1, since the segment from c

does not satisfy the “preceding” requirement (this follows from the fact that it is shorter).

So, the first n steps of MWA$^{\leftarrow}$ will result with a_1^t. We are now left with c. Applying the algorithm now to c, we get c^t. Therefore, the final result of the algorithm is

$$f^t = a_2 + c^t.$$

Now $f^t = a_2 + c_t$ implies

$$c_t = c^t.$$

This ends the proof of the claim in this case.

(2) We now proceed to the remaining case

$$C_1 \leq D_2.$$

We now analyze what MWA$^{\leftarrow}$ tells us here regarding f^t and c^t.

We illustrate f and c by the following example:



(6.33)

We apply MWA$^{\leftarrow}$ to $f = a_1 + c$. The first step starts with D_1, and further, we get points $D_1 - 1, \ldots, D_2 - 1$ (observe that $D_2 < D_1$, which follows from $A_2 + D_2 < A_1 + D_1$ and the assumption on supports). At each stage, there is only one segment with that end. The next point, D_2, is different. Here we have two segments ending with this point. One segment is $[D_2 - d + 1, D_2]$ (belonging to a_1), while the other one is obtained by (possible) linking of segments in a_2. Since D_2 is the biggest end in a_2, the length of the segment ending with D_2 in c is at least e. Therefore, in this case we have two segments ending with D_2: $[D_2 - d + 1, D_2]$ and the other one whose length is at least e.

We have three possibilities. If the segment from a_1 is shorter then the one from c, we take its end by MWA$^{\leftarrow}$ (the lexicographic requirement and “preceding” requirement are fulfilled). If the segments are of the same length, they are equal, and we take the end of the one from a_1. The last possibility is that the segment from a_1 is longer then the one from c. But then the “preceding” requirement tells us that we need to take the end of the segment from a_1, since the segment from c does not satisfy the “preceding” requirement (this follows from the fact that it is shorter).

Continuing in this way, we get the segment $[D_1 - n + 1, D_1]^{(\rho)}$, made entirely from the ends of segments from a_1.

Continuing in the above way, starting with points $D_1 - 1, \dots, D_2 + 1$, we get segments $[D_1 - n, D_1 - 1]^{(\rho)}$, $[D_1 - 1 - n, D_1 - 2]^{(\rho)}$, …, $[D_2 - n + 2, D_2 + 1]^{(\rho)}$, all from the ends of segments coming from a_1.

Now we apply MWA$^{\leftarrow}$ to the remaining segments, starting with D_2. We have two possibilities for the segment from which to take it: one is $[C_1, D_2]$ coming from a_1, and the other is $[X, D_2]$ from c (here $X \leq C_2$ depending on possible linkings). Since $C_1 < C_2$, we must start with the end of the segment coming from a_1. Therefore, we take D_2 from the segment coming from a_1, and continuing MWA$^{\leftarrow}$ in the above way, we get the segment $[D_2 - n + 1, D_2]^{(\rho)}$. This segment is obtained completely from the ends of the segments coming from a_1 in the same way as was previously explained.

In our example, we are now in the following situation:



(6.34)

We must start the next step with the end D_2 of the segment $[X, D_2]$ from c (which we have we considered above). Recall that this segment was longer than $[C_1, D_2]$. Therefore, in all further stages of MWA$^{\leftarrow}$ starting with this point, we will have segments of the same length as $[X, D_2]$, or longer. Therefore, at this step, only the segments from c will take part.

Now we go to $D_2 - 1$. Then we need to start with a segment coming from a_1, since it is shorter. In the same way as before, we now complete this step with segments coming from a_1.

We go once more, starting with $D_2 - 1$. We choose the shortest segment ending with $D_2 - 1$. There is only one possibility, and it comes from c. Its beginning is C_2 or smaller. Therefore, it is longer than the segments coming from a_1 at this step. Therefore, all the segments used at this step come from c.

We continue these steps with beginnings $D_2 - 2$, …, C_1 (possibly several times with each of them). In the same way as above, at each step we shall deal either with segments coming from a_1, or with segments coming from c (if we start with the end of a segment coming from a_1, this is clear; for the end of a segment coming from a_2, $C_2 < C_1$ implies that we complete the step with segments coming from a_2)

Further, the remaining steps take part only inside segments coming from c. The final result of the algorithm is

$$f^t = a_2 + c^t.$$

Since we have also $f^t = a_2 + c_t$ Thus

$$c_t = c^t.$$

This completes the proof of the claim in this case.

Now we finish the proof of the lemma. We know from earlier

$$c < a_2$$

and

$$c_t < a_2^t.$$

Now $c_t = c^t$ implies

$$c^t < a_2^t.$$

The preceding lemma now implies a contradiction. The proof of the lemma is now complete. □

7. Reducibility criterion

Theorem 7.1. *Let* $\pi_1 = u_{ess}(\delta_1, m_1)$ *and* $\pi_2 = u_{ess}(\delta_2, m_2)$ *for some* $\delta_1, \delta_2 \in \mathcal{D}$ *and some positive integers* m_1, m_2. *Then*

$$\pi_1 \times \pi_2$$

is reducible if and only if the underlying sets of the cuspidal supports of π_1 *and* π_2 *are linked and* π_1 *and* π_2 *are crossed.*

We now explain when the reducibility happens in a different way.

First, to have reducibility, we need to have both representations π_1 and π_2 supported on the same cuspidal $\mathbb{Z}$-line, i.e., in $\{\nu_\rho^k \rho; k \in \mathbb{Z}\}$ for some $k \in \mathcal{C}$. We assume this in what follows.

Write

$$\pi_i = L(([A_i, B_i], [A_i + 1, B_i + 1], \dots, [C_i, D_i])^{(\rho)}), \ i = 1, 2. \tag{7.35}$$

We can always chose ρ so that $A_i \in \mathbb{Z}$. Here

$$A_i \le B_i, C_i \le D_i$$

and

$$A_i + D_i = B_i + C_i$$

for $i = 1, 2$. We use the shorthand

$$\pi_i = u_{ess}\begin{pmatrix} A_i & & B_i & \\ & C_i & & D_i \end{pmatrix}^{(\rho)}$$

To have reducibility, the linking condition for the underlying sets of the cuspidal supports must be satisfied. This implies $A_1 \neq A_2$. Without lost of generality we can assume

$$A_1 < A_2. \tag{7.36}$$

Now, the linking condition is equivalent to

$$D_1 < D_2 \tag{7.37}$$

and

$$A_2 \leq D_1 + 1. \tag{7.38}$$

The crossing condition is now equivalent to

$$B_1 < B_2 \text{ and } C_1 < C_2.$$

Write

$$\begin{pmatrix} A & & B & \\ & C & & D \end{pmatrix} <_{strong} \begin{pmatrix} A' & & B' & \\ & C' & & D' \end{pmatrix}$$

if $A < A', B < B', C < C'$ and $D < D'$.

Let us summarize: If π_1 and π_2 are not supported on the same cuspidal $\mathbb{Z}$-lines, then $\pi_1 \times \pi_2$ is irreducible. If they are supported by the same cuspidal $\mathbb{Z}$-line. Then we can write them as

$$\pi_i = u_{ess}\begin{pmatrix} A_i & & B_i & \\ & C_i & & D_i \end{pmatrix}^{(\rho)}$$

where $A_i, B_i, D_i, D_i \in \mathbb{Z}$,

$$A_i \leq B_i, C_i \leq D_i$$

and

$$A_i + D_i = B_i + C_i$$

for $i = 1, 2$.

Theorem 7.2. *The representation* $\pi_1 \times \pi_2 = u_{ess}\begin{pmatrix} A_1 & & B_1 & \\ & C_1 & & D_1 \end{pmatrix}^{(\rho)} \times u_{ess}\begin{pmatrix} A_2 & & B_2 & \\ & C_2 & & D_2 \end{pmatrix}^{(\rho)}$ *reduces if and only if*

$$[A_1, D_1]_{\mathbb{Z}} \cup [A_2, D_2]_{\mathbb{Z}}$$

is a $\mathbb{Z}$*-segment, and*

$$\begin{pmatrix} A_1 & & B_1 & \\ & C_1 & & D_1 \end{pmatrix} <_{strong} \begin{pmatrix} A_2 & & B_2 & \\ & C_2 & & D_2 \end{pmatrix} \quad or \quad \begin{pmatrix} A_2 & & B_2 & \\ & C_2 & & D_2 \end{pmatrix} <_{strong} \begin{pmatrix} A_1 & & B_1 & \\ & C_1 & & D_1 \end{pmatrix}.$$

8. Relation with a result of C. Mœglin and J.-L. Waldspurger

We follow the notation of [15], and consider here the field case.

8.1. **Some notation.** Let $\delta \in \mathcal{D}^u$. As in [15], write

$$\delta[s] = \nu^s \delta.$$

Consider the following two parameters attached to δ, the (unitarizable) cuspidal representation ρ and $t \in (1/2)\mathbb{Z}_{\geq 0}$:

$$\delta = \delta([-t,t]^{(\rho)}).$$

Let $a, b \in \mathbb{R}$ such that $b - a \in \mathbb{Z}_{\geq 0}$. Let

$$\begin{aligned} J(\delta,a,b) &= L(\nu^a\delta, \nu^{a+1}\delta, \dots, \nu^b\delta) = L(\delta[a], \delta[a+1], \dots, \delta[b]) \\ &= L(\delta([-t,t]^{(\rho)})[a], \delta([-t,t]^{(\rho)})[a+1], \dots, \delta([-t,t]^{(\rho)})[b]) \\ &= L(\delta([a-t,a+t]^{(\rho)}), \delta([a+1-t,a+1+t]^{(\rho)}), \dots, \delta([b-t,b+t]^{(\rho)})). \end{aligned}$$

Therefore for $\delta = \delta(\rho, 2t+1) = \delta([-t,t]^{(\rho)})$ we have

$$J(\delta,a,b) = u_{ess}\begin{pmatrix} a-t & & a+t & \\ & b-t & & b+t \end{pmatrix}^{(\rho)}.$$

8.2. **The linking condition of** [15] **for essentially Speh representations.** Take two essentially Speh representations $J(([-t,t]^{(\rho)}),a,b)$ and $J(\delta([-t',t']^{(\rho')}),a',b')$ as above. Then they are called linked if

1. $\rho \cong \rho'$;
2. $(a-t)-(a'-t') \in \mathbb{Z}$;
3. $b > b' + |t-t'|, \quad a > a' + |t-t'|$ and $a - b' \leq 1 + t + t'$, or
   $b' > b + |t-t'|, \quad a' > a + |t-t'|$ and $a' - b \leq 1 + t + t'$.

We analyze the first condition in (3). This condition is equivalent to the fact that the following hold:

$b - t > b' - t', \quad a - t > a' - t'$ and $a - t \leq b' + t' + 1$

and

$b + t > b' + t', \quad a + t > a' + t'$ and $a - t \leq b' + t' + 1$.

Write

$$\begin{pmatrix} A & & B & \\ & C & & D \end{pmatrix} := \begin{pmatrix} a-t & & a+t & \\ & b-t & & b+t \end{pmatrix},$$

$$\begin{pmatrix} A' & & B' & \\ & C' & & D' \end{pmatrix} := \begin{pmatrix} a'-t' & & a'+t' & \\ & b'-t' & & b'+t' \end{pmatrix}.$$

Then the above condition becomes

$C' < C, \quad A' < A$ and $A \leq D' + 1$

and

$D' < D, \quad B' < B$ and $A \leq D' + 1.$

This and Lemma in I.6.3 of [15] tell us that for the field case, [15] proves one implication of the reducibility criterion that we got in this paper.

References

[1] Aubert, A. M., *Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d'un groupe réductif p- adique*, Trans. Amer. Math. Soc. 347 (1995), 2179-2189; *Erratum*, Trans. Amer. Math. Soc 348 (1996), 4687-4690

[2] Badulescu, A. I., *Global Jacquet-Langlands correspondence, multiplicity one and classification of automorphic representations (with an appendix by Neven Grbac)*, Invent. Math. 172 no. 2 (2008), 383-438.

[3] Badulescu, A. I., *On p-adic Speh representations*, Bulletin de la SMF, to appear.

[4] Badulescu, A. I., Henniart, G., Lemaire, B. and Sécherre, V., *Sur le dual unitaire de* $GL_r(D)$, Amer. J. Math. 132 no. 5 (2010), 1365-1396.

[5] Badulescu, A. I., Lapid, E. Míngues, A., *Une condition suffisante pour l'irreducibilite d'une induite parabolique de GL(m,D)*, Ann. Inst. Fourier, to appear.

[6] Badulescu, A. I. and Renard, D. A., *Sur une conjecture de Tadić*, Glasnik Mat. 39 no. 1 (2004), 49-54.

[7] Badulescu, A. I. and Renard, D. A., *Zelevinsky involution and Mœglin-Waldspurger algorithm for* $GL_n(D)$, Functional analysis IX, 9–15, Various Publ. Ser. (Aarhus), 48, Univ. Aarhus, Aarhus, 2007.

[8] Badulescu, A. I. and Renard, D. A., *Unitary dual of* GL_n *at archimedean places and global Jacquet-Langlands correspondence*, Compositio Math. 146 no. 5 (2010), 1115-1164

[9] Bernstein, J. and Zelevinsky, A.V., *Induced representations of reductive p-adic groups I* , Ann. Sci. École Norm Sup. 10 (1977), pp. 441-472.

[10] Casselman, W., *Introduction to the theory of admissible representations of p-adic reductive groups*, preprint (`http://www.math.ubc.ca/~cass/research/pdf/p-adic-book.pdf`).

[11] Deligne, P., Kazhdan, D. and Vignéras, M.-F., *Représentations des algèbres centrales simples p-adiques*, in book "Représentations des Groupes Réductifs sur un Corps Local" by Bernstein, J.-N., Deligne, P., Kazhdan, D. and Vignéras, M.-F. , Hermann, Paris, 1984.

[12] Lapid, E. and Minguez, A., *On a determinantal formula of Tadić, Amer. J. Math.*, to appear.

[13] Lapid, E., Muić, G. and Tadić, M., *On the generic unitary dual of quasisplit classical groups,* Int. Math. Res. Not. no. 26 (2004), 1335–1354.

[14] Mœglin, C. and Waldspurger, J.-L., *Sur l'involution de Zelevinski*, J. Reine Angew. Math. 372 (1986), 136-177.

[15] Mœglin, C. and Waldspurger, J.-L., *Le spectre residuel de* $GL(n)$, Ann. Sci. École Norm. Sup 22 (1989), 605-674.

[16] Muić, G. and Tadić, M., *Unramified unitary duals for split classical p–adic groups; the topology and isolated representations*, in "On Certain L-functions", Clay Mathematics Proceedings 13, 2011, 375-438.

[17] Rodier, F. *Représentations de* $GL(n,k)$ *où* k *est un corps p-adique*, Séminaire Bourbaki no 587 (1982), Astérisque 92-93 (1982), 201-218.

[18] Schneider, P. and Stuhler, U., *Representation theory and sheaves on the Bruhat-Tits building*, Publ. Math. IHES 85 (1997), 97-191.

[19] Sécherre, V., *Proof of the Tadić conjecture (U0) on the unitary dual of $GL_m(D)$*, J. reine angew. Math. 626 (2009), 187-203.

[20] Speh, B., *Unitary representations of $GL(n,\mathbb{R})$ with non-trivial (g,K)- cohomology*, Invent. Math. 71 (1983), 443-465.

[21] Tadić, M., *Classification of unitary representations in irreducible representations of general linear group (non-archimedean case)*, Ann. Sci. Ecole Norm. Sup. 19 (1986), 335-382.

[22] Tadić, M., *Topology of unitary dual of non-Archimedean $GL(n)$,* Duke Math. J. 55 no. 2 (1987), 385-422.

[23] Tadić, M., *Induced representations of $GL(n,A)$ for p-adic division algebras A*, J. Reine Angew. Math. 405 (1990), 48-77.

[24] Tadić, M., *Structure arising from induction and Jacquet modules of representations of classical p-adic groups*, J. of Algebra 177 (1995), 1-33.

[25] Tadić, M., *On reducibility of parabolic induction,* Israel J. Math. 107 (1998), 29-91.

[26] Tadić, M., *On the representation theory of $GL(n)$ over a p-adic division algebra and unitarity in the Jacquet-Langlands correspondences*, Pacific J. Math. 223 (2006), 167-200.

[27] Tadić, M., *$GL(n,\mathbb{C})\hat{}$ and $GL(n,\mathbb{R})\hat{}$*, in "Automorphic Forms and L-functions II, Local Aspects", Contemporary Mathematics 489, 2009, 285-313.

[28] Tadić, M., *On reducibility and unitarizability for classical p-adic groups, some general results*, Canad. J. Math. 61 (2009), 427-450.

[29] Tadić, M., *On interactions between harmonic analysis and the theory of automorphic forms*, preprint (`http://www.hazu.hr/~tadic/50-harmonic-auto.pdf`).

[30] Zelevinsky, A. V., *Induced representations of reductive p-adic groups II. On irreducible representations of GL(n)*, Ann. Sci. École Norm. Sup. 13 (1980), pp. 165-210.

Department of Mathematics, University of Zagreb, Bijenička 30, 10000 Zagreb, Croatia,
Email: `tadic@math.hr`